

# Australian College of Midwives Draft Position Statement on Bed-sharing and Co-sleeping

## Consultation Paper: 20<sup>th</sup> December 2013

The Australian College of Midwives (ACM) is currently developing a Position Statement on bed-sharing and co-sleeping. This Draft document has been developed by the ACM Scientific Research Advisory Committee (SRAC) based on available evidence. It is now ready for wider consultation and consideration by ACM membership and stakeholders. Your feedback is very important because it supports the ACM to develop its Position Statement based on the latest evidence. Once the consultation period ends, the feedback will be collated and changes will be considered. The final draft will be reviewed once more by SRAC, endorsed by the ACM Board and then published on the ACM website (<http://www.midwives.org.au>).

The Draft document is now ready for wider consultation and consideration by ACM membership and stakeholders. Your feedback is very important because it supports the ACM to develop its Position Statement based on best practice as well as professional and regulatory codes, standards and frameworks. Once the consultation period ends, the feedback will be collated and changes will be considered. The final draft will be endorsed by the ACM Board and then published on the ACM website (<http://www.midwives.org.au>).

Submissions will be published on the ACM website to encourage discussion and inform the community. We will not publish any material provided in-confidence. Submissions will be published as soon as possible after the end of the public comment period.

***DEADLINE FOR SUBMISSIONS to ACCYPN for collated Group Response:  
COB Monday 10 February 2014***

Submissions received after this date will not be considered.

### ***HOW TO SUBMIT for ACCYPN Collated Response:***

*Please respond with comments via email direct to the ACCYPN Secretariat:  
[info@accypn.org.au](mailto:info@accypn.org.au)*

*A Group Response on behalf of ACCYPN will be collated and forwarded direct to  
Australian College of Midwives*

*Please head your submission: Consultation - Bed-sharing and Co-sleeping. Also, give your  
name, job title, and organisation where you are employed.*

## ACM Draft Position Statement on Bed-sharing and Co-sleeping

The ACM supports the facilitation of women and their families making informed choices in maternity and child care, including decisions about where their baby sleeps.

The ACM advocates a risk minimisation approach in which the individual family's circumstances are considered when providing advice about infant care practices parents will use in caring for their baby. The ACM believes that in ensuring the safety of babies, it is crucial to respect and support cultural norms and practices.

The ACM does not support a risk elimination approach in which all parents are informed not to bed-share or co-sleep with their baby under any circumstances.

### Definitions

There are various definitions that have been used in the literature to describe the practices of bed-sharing and co-sleeping<sup>1</sup>.

The definitions used for co-sleeping and bed-sharing in this position statement are consistent with definitions used in safe sleeping guidelines nationally<sup>2,3</sup> and internationally<sup>1,4</sup> by health professionals and policy development, and allow for differentiation of the risks associated with solitary sleeping, room sharing and environments in which baby and caregiver share the same sleep surface.

*Bed-sharing refers to bringing a baby onto a sleep surface when co-sleeping is possible, whether intended or not<sup>1,3-5</sup>.*

*Co-sleeping is defined as a mother and/or her partner (or any other person) being asleep on the same sleep surface as the baby<sup>1,3,4</sup>.*

The ACM identifies 12 key principles for midwives who care for mothers and their babies and who provide parent education and advice relating to safe infant sleeping, including bed-sharing and co-sleeping practices.

### Key Take Home Messages

- Bed-sharing and co-sleeping are common, valued infant care practices in both developed and developing countries across the world.
- Bed-sharing and co-sleeping are associated with many benefits however the risk of sudden unexpected death in infancy including SIDS and fatal sleeping accidents is increased in certain circumstances.
- Bed-sharing and breastfeeding are integrated infant care practices: breastfeeding promotes bed-sharing, while bed-sharing promotes breastfeeding.
- Many parents will bed-share with their baby, whether they intend to, or not.
- All parents should be provided with clear and impartial information about the benefits and contraindications to sharing a sleep surface using risk minimisation guidelines in order to make informed decisions.

**How do you feel about the opening statements?**

**How do you feel about the Definitions?**

**How do you feel about the Key Take Home Messages?**

## Key principles

1. Co-sleeping and bed-sharing between mothers and their babies is an established and valued practice in most cultures; and considered the social norm for approximately 90% of the world's population<sup>6-10</sup>. Sharing a sleep surface with a baby is a common infant care practice in Australia, with rates reported between 51-80% depending on infant age at time of measurement<sup>5,11</sup>.
2. Bed-sharing and co-sleeping with a baby has been associated with many benefits<sup>12,13</sup>. Benefits include enhanced maternal-infant bonding and maternal responsiveness<sup>8,14-17</sup>; improved settling with reduced crying<sup>15</sup>; improved maternal and infant sleep and increased arousals<sup>4,15,18,19</sup>; increased duration of breastfeeding<sup>8,20-22</sup>; and reduced formula supplementation<sup>16,22,23</sup>. Longitudinal studies have suggested that those who shared the parental bed as babies and children become adults with higher self-esteem, and better social skills and emotional outcomes<sup>8,24,25</sup>.
3. Bed-sharing and co-sleeping have been associated with an increased risk of sudden unexpected death in infancy (SUDI), including Sudden Infant Death Syndrome (SIDS) and fatal sleeping accidents, in certain circumstances<sup>12,17,26-28</sup>. Please see Glossary for definitions used for SUDI, SIDS and fatal sleeping accidents.
4. The risk associated with shared sleep surfaces is significantly increased by the presence of other known risk factors for sudden infant death including antenatal and postnatal exposure to tobacco smoking, prone sleep position, parental drug (prescribed or illicit) and alcohol consumption, soft sleep surfaces, multiple bed-sharers, maternal sedation and obesity. Sleeping on a sofa with a baby significantly increases the risk and should be avoided. Babies most at risk of SUDI are those born preterm, of low birth weight, and babies less than 11 weeks of age<sup>27-34</sup>.
5. The *Sleep Safe, My Baby* public health recommendations are underpinned by evidence which demonstrates that avoidance of potentially modifiable risk factors will reduce the risk of sudden unexpected deaths in infancy for vulnerable infants. See Appendix A.
6. Room-sharing is associated with a reduced risk of SUDI. Parents should be encouraged to keep their babies in their bedroom at night for the first six months of life<sup>35,36</sup>.
7. Breastfeeding is protective against SUDI<sup>37</sup>. It is desirable to encourage and support exclusive breastfeeding; and bed-sharing is an infant care practice significantly associated with breastfeeding longevity<sup>10</sup>. Breastfeeding and sharing a sleep

surface constitute an integrated care system which is mutually reinforcing; breastfeeding promotes shared sleep which increases breastfeeding frequency and extends duration of breastfeeding in months<sup>8,21,22,38-40</sup>.

8. Many parents will bring their baby into bed as this helps them to care for their baby at night, particularly if they are breastfeeding. Many parents will fall asleep with their baby, whether they intend to, or not<sup>1,17,40-44</sup>.
9. The ACM supports the facilitation of women and their families making informed choices in maternity and child care. Informed decision making, informed consent and right of refusal are accepted legal principles in Australia. Each and every woman and her family have the right to make informed decisions, including consent and refusal of any aspect of her care. Women and their families must be respected in the choices that they make.
10. Midwives have a primary responsibility to ensure that their decisions, recommendations and practices are focused on the individual needs and safety of the woman and her baby/babies.
11. Evidence has shown that health professionals are in a unique position to influence the behaviour of parents and caregivers, by modelling safe infant sleep practices in hospital and providing parents with information and support strategies, to ensure parent practices used at home are consistent with public health safe infant sleeping recommendations and risk minimisation strategies in shared sleeping environments<sup>11,22,33,34,44,45</sup>. See Appendix B for risk minimisation strategies to reduce the risk of SUDI associated with shared sleeping in the home environment.
12. Compliance with safe sleeping recommendations and risk minimisation strategies for shared sleeping environments is reliant on the ability of midwives to engage parents, identify individual sleeping environments and provide evidence-based advice to parents that aims to reduce risk associated with all sleeping environments, particularly in circumstances where parent-infant sleep on the same sleep surface is likely to occur<sup>5</sup>.

**How do you feel about the key principles?**

**What changes would you suggest to the key principles?**

### **Achieving best practice**

To achieve best practice in ensuring parents are provided with unbiased evidence-based information in order to make an informed decision about the infant care practices they will use in caring for their baby, it is necessary for consumers (e.g. Australian Breastfeeding Association, parent consumers), professional colleges, support organisations (e.g. SIDS and Kids), education providers, health systems, Australian and State and Territory governments and policy makers to work together to:

- Ensure that midwifery practice is underpinned by national evidence-based safe infant sleeping recommendations. The *SIDS and Kids Safe Sleep My Baby* Australian public health campaign to reduce the risk of sudden unexpected death in infancy,

including SIDS and fatal sleeping accidents<sup>36</sup> supports a risk minimisation approach to bed-sharing and co-sleeping. See Appendix A;

- Promote consistency of evidence-based information to parents about the benefits, contraindications and strategies to enhance safety associated with bed-sharing and co-sleeping environments so that parents and carers can make informed decisions regarding sleeping arrangements for their baby in the hospital and/or in the home environment;
- Provide parents with clear and impartial information on the benefits of, and contraindications to, sharing a sleep surface using risk minimisation guidelines<sup>3,9,10</sup>;
- Discuss the benefits and contra-indications to bed-sharing and co-sleeping with all mothers in the antenatal period (by 36 weeks gestation) and again in the early post-natal period up to 12 months of age to allow them to make a fully informed choice. This should include written information about safe infant sleeping and risk minimisation strategies associated with bed-sharing and co-sleeping in a format appropriate to their language needs. See Appendix B;
- Develop local evidence-based guidelines on mother-baby bed-sharing and co-sleeping for in-patient facilities including postnatal, residential care and paediatric tertiary facilities where families with young infants are cared for, in consultation with consumers;
- Implement a risk assessment of mothers and babies in inpatient facilities prior to mother taking baby into bed for feeding and/or settling which considers the clinical condition of both mother and baby, the safety of the physical environment, and the level of supervision required. Assessment should identify risks and risk minimisation strategies, and specifically address circumstances where co-sleeping is not recommended. Circumstances for mothers and babies may quickly change, therefore this risk assessment will need to be reviewed as required;
- Consider cultural, social and family circumstances which may impact on a caregiver's choice of infant sleep location and provide appropriate education and information relating to these environments to facilitate informed decision-making by parents;
- Provide educational opportunities to ensure that midwives maintain their clinical skills and knowledge of evidence-based parenting advice in co-sleeping and bed-sharing practice so that they are competent and confident to provide care for women and babies and discuss the issues objectively with women to enable informed decision-making;
- Undertake research into co-sleeping and bed-sharing practices. There is limited Australian research on the prevalence and incidence of bed-sharing and co-sleeping in contemporary Western societies and Indigenous communities using current definitions which differentiate between shared sleeping environments. Further research examining the unique nature of the postnatal maternity

environment, factors which influence health professional current clinical practice, and the impact of health professional modelling in relation to ongoing maternal co-sleeping and bed-sharing behaviours is required<sup>46</sup>.

**How do you feel about the recommendations for achieving best practice?**

**What changes would you suggest to the recommendations for achieving best practice?**

### Resources to guide practice

This position statement has been informed by a literature review which can be found at <http://www.midwives.org.au/scripts/cgiip.exe/WService=MIDW/ccms.r?pageid=10209>

The ACM supports the use of the following resources to guide midwives in their practice:

- Clinical Skills Development Centre (2013) Safe Infant Sleeping (SUDI) eLearning Program. Collaboration between SIDS and Kids and QH Clinical Skills Development Centre. (Author: Professor Jeanine Young) Endorsed by Australian College of Midwives and Australian College of Nurses [5 Continuing Education Points]. Available <http://www.sdc.qld.edu.au/courses/103>
- Clinical Skills Development Centre (2013) Aboriginal and Torres Strait Islander Safe Infant Sleeping (ISIS) eLearning Program. (Author: Professor Jeanine Young & Leanne Craigie) Available <http://www.sdc.qld.edu.au/courses/123>
- UNICEF UK Baby Friendly Initiative. (2012). Caring For Your Baby at Night: A guide for parents. In UNICEF UK Baby Friendly Initiative (Ed.). London: UNICEF.
- Blair P, Inch S. (2012) The health professional's guide to "Caring for your Baby at night". In UNICEF UK Baby Friendly Initiative (Ed.). London: UNICEF.
- UNICEF Leaflet 'Sharing a bed with your baby: A guide for breastfeeding mothers' (May 2008) available at [www.babyfriendly.org.uk/pdfs/sharingbedleaflet.pdf](http://www.babyfriendly.org.uk/pdfs/sharingbedleaflet.pdf).
- UNICEF UK Baby Friendly Initiative. (2004). *Babies sharing their mothers' bed while in hospital: A sample policy*. London: UNICEF UK Baby Friendly Initiative.
- SIDS and Kids (2012) Safe Sleeping: A guide to assist in sleeping your baby safely. Melbourne: SIDS and Kids. [www.sidsandkids.org](http://www.sidsandkids.org)
- SIDS and Kids (2013) Information statement: *Sleeping with a baby*. First published September, 2007. Canberra: SIDS and Kids. <http://www.sidsandkids.org> (Being updated in 2013 by December 2013).
- Young J, Watson K, Ellis L, Raven L. (2012) Responding to the evidence: Breastfeed baby if you can – the sixth public health recommendation to reduce the risk of sudden unexpected death in infancy. *Breastfeeding Review* 20(1): 7-15.
- Infant Sleep Information Source (ISIS): Information for Parents and Carers Collaboration between Durham University Parent-Infant Sleep Laboratory, La Leche League, National Childbirth Trust and UNICEF UK Baby Friendly Initiative funded by the Economic and Social Research Council (ESRC). <http://www.isisonline.org.uk/>

- NCT *Position statement: co-sleeping and bed-sharing*. Available from: <http://www.nct.org.uk/press-office/positionstatements/transitionparenthood>
- Royal College of Midwives (2004) *Bed-sharing and co-sleeping: Position Statement No. 8*
- Queensland Health. (2008) *Safe infant care to reduce the risk of sudden unexpected deaths in infancy: Policy statement and minimum practice standards*. Brisbane: Maternity, Child Health and Safety Branch, Queensland Health. [ISBN: 9781921447280] Available <http://www.sidsandkids.org/wp-content/uploads/Safe-Infant-Sleeping-Policy-Statement-and-Guidelines.pdf>
- Queensland Health (2013) *Safe Infant Sleeping Policy, Education and Resources*. Available <http://www.health.qld.gov.au/child-youth/webpages/safesleeping.asp>
- Australian Breastfeeding Association (2013) *Breastfeeding, Co-sleeping and Sudden Unexpected Deaths in Infancy*. (Authors Kam R, Young J). Available from [www.breastfeeding.asn.au/bfinfo/breastfeeding-co-sleeping-and-sudden-unexpected-deaths-infancy](http://www.breastfeeding.asn.au/bfinfo/breastfeeding-co-sleeping-and-sudden-unexpected-deaths-infancy)
- The Academy of Breastfeeding Medicine Protocol Committee (2008) *ABM Clinical Protocol #6: Guideline on Co-sleeping and Breastfeeding*. Revision, March 2008. *Breastfeeding Medicine* 3(1): 38-43. Queensland Health's Child Health Information Fact sheet entitled *Safe Sleeping for babies: Reducing the risk of sudden infant death* is available for distribution to parents and staff at [www.health.qld.gov.au/publications/childhealth](http://www.health.qld.gov.au/publications/childhealth) which addresses strategies to reduce risk associated with shared sleep environments

**How do you feel about the resources to guide midwives in their practice?**

**What changes would you suggest to the resources to guide midwives in their practice?**

## Glossary

### Definitions of terms used in this position statement and supporting documents

Term	Definition / Explanation / Details	Source
<b>Bed-sharing</b>	Bed-sharing refers to bringing baby onto a sleep surface when co-sleeping is possible, whether intended or not.	UNICEF Leaflet "Sharing a bed with your baby: A guide for breastfeeding mothers" (May 2008) available at <a href="http://www.babyfriendly.org.uk/pdfs/sharingbedleaflet.pdf">www.babyfriendly.org.uk/pdfs/sharingbedleaflet.pdf</a> UNICEF U.K. Baby Friendly Initiative (2004). <i>Babies sharing their mothers' bed while in hospital: A sample policy</i> . UNICEF UK Baby Friendly Initiative, London. Infant Sleep Information Source (2013) Definitions of terms used on this site. Available 18 October 2013 from <a href="http://www.isisonline.org.uk/hcp/research_evidence/definitions/">http://www.isisonline.org.uk/hcp/research_evidence/definitions/</a>
<b>Co-sleeping</b>	Co-sleeping is defined as a mother and /or her partner (or any other person) being asleep on the same sleep surface as the baby.	UNICEF Leaflet "Sharing a bed with your baby: A guide for breastfeeding mothers" (May 2008) available at <a href="http://www.babyfriendly.org.uk/pdfs/sharingbedleaflet.pdf">www.babyfriendly.org.uk/pdfs/sharingbedleaflet.pdf</a> UNICEF U.K. Baby Friendly Initiative (2004). <i>Babies sharing their mothers' bed while in hospital: A sample policy</i> . UNICEF UK Baby Friendly Initiative, London. Infant Sleep Information Source (2013) Definitions of terms used on this site. Available 18 October 2013 from <a href="http://www.isisonline.org.uk/hcp/research_evidence/definitions/">http://www.isisonline.org.uk/hcp/research_evidence/definitions/</a>
<b>Sharing the same sleep surface</b>	Includes the practices of <i>bed-sharing</i> and <i>co-sleeping on the same sleep surface</i> . This terminology allows differentiation of the risks associated with solitary sleeping (baby sleeping in a separate room), room-sharing, and environments in which the baby and caregiver share the same sleep surface	SIDS and Kids (2013) Information Statement: <i>Sleeping with a baby</i> . Original .September, 2007. Melbourne: SIDS and Kids. Downloaded 18 <sup>th</sup> October 2013 from <a href="http://www.sidsandkids.org.au">http://www.sidsandkids.org.au</a> Queensland Health. (2008) Safe infant care to reduce the risk of sudden unexpected deaths in infancy: Policy statement and minimum practice standards. Brisbane: Maternity, Child Health and Safety Branch, Queensland Health. [ISBN: 9781921447280] <a href="http://www.sidsandkids.org/wp-content/uploads/Safe-Infant-Sleeping-Policy-Statement-and-Guidelines.pdf">http://www.sidsandkids.org/wp-content/uploads/Safe-Infant-Sleeping-Policy-Statement-and-Guidelines.pdf</a>
<b>SUDI</b>	SUDI is defined as the "sudden, unexpected death of an infant usually occurring during sleep, in which the cause is not immediately obvious". SUDI is essentially a research classification, and refers to a broad category of sudden and unexpected deaths which include Sudden Infant Death Syndrome (SIDS), infections or anatomical or developmental abnormalities not recognised before death, fatal sleep accidents due to unsafe sleep environments, and sudden unexpected deaths that are revealed by investigations to have been the result of non-accidental injuries.	The State of Queensland (Commission for Children and Young People and Child Guardian Queensland) (2012) <i>Annual Report: Deaths of children and young people, Queensland 2011-2012</i> . Commission for Children and Young People and Child Guardian Queensland, Brisbane.  Sidebotham P, Bajanowski T, Keens T, Kenner T, Kerbl R, Kurz R, Mitchell EA, Moon R, Taylor B, Venneman M, Young J, Zotter H. (2010) Proposal for an international classification of SUDI: A response to Blair, Byard and Fleming. <i>Scandinavian Journal of Forensic Science</i> 16(1): 9-11.
<b>SIDS</b>	Sudden Infant Death Syndrome (SIDS) is defined as "the sudden and unexpected death of an infant under one year of age, with onset of the lethal episode apparently occurring during sleep, that remains unexplained after a thorough investigation including performance of a complete autopsy and review of the	The State of Queensland (Commission for Children and Young People and Child Guardian Queensland) (2012) <i>Annual Report: Deaths of children and young people, Queensland 2011-2012</i> . Commission for Children and Young People and Child Guardian Queensland, Brisbane.  Krous H, Beckwith J, Byard R, Bajanowski T, Corey T, Cutz E, Hanzlick R, Keens T, Mitchell, E. (2004) Sudden infant death syndrome and unclassified infant deaths: a definitional and diagnostic approach. <i>Pediatrics</i> 114(1): 234-238.



	circumstances of death and the clinical history.' 'SIDS is a subset of SUDI and is a classification of exclusion.	
--	---	--

Term	Definition / Explanation / Details	Source
<b>Fatal sleeping accident</b>	A death occurring during sleep, as a result of an accident, such as a fall, or suffocation, or mechanical asphyxiation. Fatal sleeping accidents are explained deaths that meet SUDI criteria.	Commission for Children and Young People and Child Guardian Queensland (2012) <i>Annual Report: Deaths of children and young people, Queensland 2011-12</i> . Commission for Children and Young People and Child Guardian Queensland, Brisbane.
<b>Passive smoking</b>	Refers to breathing tobacco smoke in the environment.	Centre for Community Child Health (2006) <i>Preventing Passive Smoking Effects on Children: Practice Resource</i> . Victorian Government: Centre for Community Child Health. <a href="http://www.rch.org.au/ccch">www.rch.org.au/ccch</a>
<b>Environmental tobacco smoke (ETS)</b>	Refers to smoke from the end of a lit cigarette or breathed out by a smoker.	Centre for Community Child Health (2006) <i>Preventing Passive Smoking Effects on Children: Practice Resource</i> . Victorian Government: Centre for Community Child Health.
<b>Room sharing</b>	Refers to sleeping the baby in a cot or other separate sleeping surface in the same room as the parents.	SIDS and Kids (2008) Information Statement: <i>Room-sharing</i> . September, 2008. Melbourne: SIDS and Kids. <a href="http://www.sidsandkids.org.au">http://www.sidsandkids.org.au</a>

**How do you feel about the Glossary?**

**What changes would you suggest to the Glossary?**

**Any other comments?**

## Appendix A

### Safe Sleep, My Baby Public Health Campaign 2012

In Australia, SIDS and Kids launched the revised national public health campaign “Sleep Safe, My Baby” in May 2012 with six key messages<sup>35,43</sup>:

1. Sleep baby on the back from birth, not on the tummy or side
2. Sleep baby with head and face uncovered
3. Keep baby smoke free before birth and after
4. Provide a safe sleeping environment night and day
5. Sleep baby in their own safe sleeping place in the same room as an adult caregiver for the first six to twelve months
6. Breastfeed baby

SIDS and Kids promotes a separate sleeping space in the same room as a caregiver for babies as the safest place for a baby to reduce the risk of SUDI, however acknowledges the many benefits of bed-sharing and co-sleeping including the support of breastfeeding. SIDS and Kids supports a risk minimisation approach that supports informed decision making by parents and provides evidence-based advice for parents who share a sleep surface with their baby<sup>2,36</sup>. These strategies are consistent with those listed in Appendix B.

## Appendix B

### Risk Minimisation Approach for Home Environments:

#### **Suggested strategies to reduce the risk of sudden unexpected deaths in infancy, including SIDS and fatal sleeping accidents, associated with co-sleeping and bed-sharing environments once at home**

In consideration of the many documented benefits of shared sleeping, the need to promote and support breastfeeding, the high prevalence of shared sleep environments in contemporary Australian society and the right of parents to make informed choices about their baby's care, the Australian College of Midwives supports the recommendation that parents should be provided with information that includes benefits, risks and strategies to reduce the risk and increase safety associated with shared sleep environments, should they decide or have no option but to share a sleep surface with their baby<sup>15,16,27,38,40</sup>.

This recommendation is consistent with, and supported by, recommendations for health professional practice proposed by UNICEF<sup>1,9,10</sup>, the Royal College of Midwives<sup>47</sup> and SIDS and Kids<sup>2,36</sup>.

In order to provide parents with information to allow informed choices about safe sleeping practices relating to shared sleep environments, parent education should include the following information:

It is *not* safe to share a sleep surface with a baby if:

- Either parent is a smoker
- Either parent is under the influence of alcohol or illicit drugs
- Either parent is under the influence of medication that causes sedation, is excessively tired, or obese.

If parents choose to share a sleeping surface with their baby, the following strategies have been demonstrated to reduce the risk of sudden unexpected deaths in infancy, including SIDS and fatal sleeping accidents<sup>1,2,5,36,48</sup>:

- Sleep baby on the back from birth – never on the tummy or side.
- If baby lies on his or her side to breastfeed, baby should be returned to the supine (back) position for sleep.
- Make sure the mattress is firm and flat (not tilted or elevated)
- Make sure that bedding cannot cover baby's face or overheat baby (use lightweight blankets and remove pillows, doonas and other soft items from the environment that could cover baby);
- Sleep baby beside one parent only, rather than between two parents, to reduce the likelihood of baby becoming covered by adult bedding (unless sleep enabler that provides for a separate sleep surface and infant bedding is being used)
- Ensure partner knows baby is in the bed.
- As an alternative to bedding, an infant sleeping bag may be used so that the baby does not share the adult bedding. (N.B. Do not *wrap* baby if sharing a sleep surface as this restricts arm and leg movement).

- Make sure baby cannot fall off the bed. A safer alternative is to place the mattress on the floor (be aware of potential situations where baby can become trapped). Do not place pillows at the side of the baby to prevent rolling off.
- Pushing the bed up against the wall can be hazardous. Babies have died after being trapped between the bed and the wall.
- Never place a baby to sleep in a bed with other children or pets (see SIDS and Kids Frequently Asked Questions for specific advice about the safest way to sleep twins at [www.sidsandkids.org](http://www.sidsandkids.org))
- **Babies must never be left alone on an adult bed or put to sleep on a sofa, bean bag, waterbed or soft, sagging mattress.**
- Three sided-cots (a cot with one side down) may be available for purchase that can be attached to the side of the bed at the same level so that the baby has a separate environment but is still in contact with a parent during sleep. *Please note that currently there is no Australian standard for three-sided cots.*

## References

1. Infant Sleep Information Source (ISIS): Information for Parents and Carers Collaboration between Durham University Parent-Infant Sleep Laboratory, La Leche League, National Childbirth Trust and UNICEF UK Baby Friendly Initiative funded by the Economic and Social Research Council (ESRC). c2013 [updated August 2013; cited November 2013] Available from: <http://www.isisonline.org.uk/>
2. SIDS and Kids Information statement: *Sleeping with a baby*. Canberra: SIDS and Kids c2006 [Being updated December 2013; cited November 2013] Available from: <http://www.sidsandkids.org>
3. Queensland Health. Safe Infant Sleeping, Co-sleeping and Bed-sharing: Guideline for Safe Infant Sleeping, Co-sleeping and Bed-sharing. Version 1.0 c2013 [Updated July 2013: cited November 2013]. QH-GDL-362: 2013.
4. UNICEF UK Baby Friendly Initiative. *Babies sharing their mothers' bed while in hospital: A sample policy*. London: UNICEF UK Baby Friendly Initiative, 2004.
5. Queensland Health. Safe infant care to reduce the risk of sudden unexpected deaths in infancy: Policy statement and minimum practice standards. Brisbane: Maternity, Child Health and Safety Branch, Queensland Health. c2008 [Written November 2008: cited November 2013] Acknowledged Author: Dr Jeanine Young [ISBN: 9781921447280] Available from: <http://www.sidsandkids.org/wp-content/uploads/Safe-Infant-Sleeping-Policy-Statement-and-Guidelines.pdf>
6. Tipene-Leach D, Abel S, Finau S, Park J, Lenna M. Maori infant care practices: implications for health messages, infant care services and SIDS prevention in Maori communities. *Pac Health Dialog* 2000; 7(1): 29-37.
7. Abel S, Park J, Tipene-Leach D, Finau S & Lennan M. Infant care practices in New Zealand: a cross cultural qualitative study. *Soc Sci Med* 2001; 53(9): 1135-1148.
8. McKenna J & McDade T. Why babies should never sleep alone: A review of the co-sleeping controversy in relation to SIDS, bedsharing and breastfeeding. *Paediatr Respir Rev* 2005; 6(2): 134-152.
9. Blair P, Inch S. The health professional's guide to "Caring for your Baby at night". In UNICEF UK Baby Friendly Initiative (Ed.). London: UNICEF, 2012.
10. UNICEF UK Baby Friendly Initiative. *Caring For Your Baby at Night: A guide for parents*. In UNICEF UK Baby Friendly Initiative (Ed.). London: UNICEF, 2012.
11. Young J, Battistutta D, O'Rourke P, Thompson JMD. *Infant Care Practices Related to Sudden Infant Death Syndrome in Queensland 2002*. Brisbane: Queensland Health, 2008.
12. Sobralcke MC, Gruber ME. Risks and benefits of parent/child bed sharing. *J Am Acad of Nurse Pract*, 2009; 21(9): 474-479. Available from: DOI: 10.1111/j.1745-7599.2009.00430
13. Morgan B, Horn A, Bergman N. Should neonates sleep alone? *Biol Psychiatr* 2011; 70(9): 817 - 825. Available from: DOI: 10.1016/j.biopsych.2011.06.018. Epub 2011 Jul 29.
14. Young, J. Bedsharing with babies: the facts. *RCM Midwives Journal* 1998; 1(11): 338-341.
15. Young J. Night-time behaviour and interactions between mothers and their infants of low risk for SIDS: a longitudinal study of room-sharing and bed-sharing. PhD, University of Bristol (United Kingdom), Bristol, 1999.

16. Baddock S. Co-sleeping: an ecological parenting practice. In Davies, L., Daellenbach R. & Kensington M (eds). *Sustainability, Midwifery and Birth*, Routledge, London, 2011; pg 207-217.
17. Ball H, Moya E, Fairley L, Westman J, Oddie S, Wright J. Infant care practices related to sudden infant death syndrome in South Asian and White British families in the UK. *Paediatr Perinat Epidemiol* 2012; 26(1):3-12. Available from: doi: 10.1111/j.1365-3016.2011.01217.x. Epub 2011 Aug 18.
18. Mosko S, Richard C, McKenna J. Infant arousals during mother-infant bed sharing: implications for infant sleep and sudden infant death syndrome research. *Pediatrics* 1997; 100(5): 841-849.
19. Mosko S, Richard C, McKenna J. Maternal sleep and arousals during bed-sharing with infants. *Sleep* 1997b; 20(2): 142-150.
20. McKenna J, Mosko S, Richard, C. Bedsharing promotes breastfeeding. *Pediatrics* 1997; 100(2 Pt 1): 214-9
21. Huang I, Hauck FR, Signore C, Airong Y, Tonse NKR, Huang TTK, Fein SB. Influence of bedsharing activity on breastfeeding duration among US mothers. *JAMA Pediatr*. [Published online September 2013; cited November 2013]. Available from: doi.10.1001/jamapediatrics.2013.2632.
22. Colson ER, Willinger M, Rybin D, Heeren T, Smith LA, Lister G, Corwin MJ. Trends and Factors Associated with Infant Bed-Sharing 1992-2010. *JAMA Pediatr*. [Published online September 2013; cited November 2013]. Available from: doi:10.1001/jamapediatrics.2013.2560.
23. Pemberton, D. Breastfeeding, co-sleeping and the prevention of SIDS. *Br J Midwifery* 2005; 13(1): 12-18.
24. Mosenkis J. The effects of childhood co-sleeping on later life development. MSc, The University of Chicago, 1998.
25. Okami P, Weisner T, Olmstead R. Outcome correlates of parent-child bedsharing: an eighteen-year longitudinal study. *J Dev Behav Pediatr* 2002; 23(4): 244-253.
26. Blair PS, Fleming PJ, Smith IJ, Ward Platt M, Young J, Nadin P, Berry PJ, Golding J and the CESDI SUDI Research Group. Babies sleeping with parents: case-control study of factors influencing the risk of the sudden infant death syndrome. *Br Med J* 1999; 319(7223): 1457-1462.
27. Blair PS, Sidebotham P, Evason-Coombe C, Edmonds M, Heckstall-Smith EMA, Fleming, P. Hazardous cosleeping environments and risk factors amenable to change: case-control study of SIDS in south-west England. *Br Med J* 2009; 339(b3666). Available from doi:http://dx.doi.org/10.1136/bmj.b3666.
28. Carpenter R, McGarvey C, Mitchell, EA, Tapping DM, Vennemann MM, Smuk S, Carpenter JR. Bedsharing when parents do not smoke: is there a risk of SIDS? AN individual level analysis of five major case-control studies. *Br Med J[Open]* 2013; 3:e002299. Available from: doi:10.1136/bmjopen-002299.
29. Blair PS, Sidebotham P, Berry PJ, Evans M, Fleming PJ. Major epidemiological changes in sudden infant death syndrome: a 20 year population-based study in the UK. *Lancet* 2006; 367(9507): 314-319.
30. Trachtenberg FL, Haas EA, Kinney HC, Stanley C, Krous HF. Risk factor changes for sudden infant death syndrome after initiation of back-to-sleep campaign. *Pediatrics* 2012; 129(4): 630-8. Available from doi:10.1542/peds.2011-1419.

31. Fleming PJ, Blair PS, Bacon C, Berry J. (Eds) Sudden unexpected deaths in infancy: the CESDI SUDI studies 1993-1996. London: The Stationery Office, 2000.
32. Commission for Children and Young People and Child Guardian [CCYPCG] Queensland, Annual Report: Deaths of children and young people, Queensland 2011-2012. Commission for Children and Young People and Child Guardian, Queensland, Brisbane, 2012.
33. NSW Child Death Review Team. Sudden Unexpected Deaths in Infancy: the New South Wales experience. Sydney: NSW Commission for Children and Young People, NSW Child Death Review Team, 2005.
34. NSW Child Death Review Team. NSW Child Death Review Team- Annual Report 2011. Sydney: NSW Ombudsman, 2012.
35. Mitchell E, Freemantle J, Young J, Byard R. Scientific consensus forum to review the evidence underpinning the recommendations of the Australian SIDS and Kids Safe Sleeping Health Promotion Programme - October 2010. *J Paediatr Child Health* 2012; 48(8): 626-633. Available from: doi:10.1111/j.1440-1754.2011.02215.x
36. SIDS and Kids. Safe Sleeping: A guide to assist in sleeping your baby safely. Melbourne: SIDS and Kids, 2012. Available from: [www.sidsandkids.org](http://www.sidsandkids.org)
37. Hauck F, Thompson J, Tanabe K, Moon R & Vennemann M. Breastfeeding and reduced risk of sudden infant death syndrome: A meta-analysis. *Pediatrics* 2012; 128(1): 103-110.
38. McKenna JJ, Ball HL, Gettler LT. Mother-infant co-sleeping, breastfeeding and sudden infant death syndrome: what biological anthropology has discovered about normal infant sleep and pediatric sleep medicine. *Am J Phys Anthropol* 2007; 134(Supplement 45):133-61.
39. Blair P, Heron J, & Fleming P. Relationship between bed sharing and breastfeeding: Longitudinal, population based analysis. *Pediatrics* 2010; 126(5): e1119-1126.
40. Ball H. The latest on bed sharing and breastfeeding. *Community Pract* 2012; 85(1): 29-31.
41. Young J, Thompson J. Recommendations for real life: the nature of shared sleep environments in Queensland and implication for effective safe infant sleeping messages. *Forensic Sci Med Pathol*, 2009; 5(2):115.
42. Dodd J. Evaluation of the Department of Health Western Australian Operational Directive Statewide Co-sleeping / Bed-sharing Policy for WA Health Hospitals and Health Services. Collaboration for Applied Research and Evaluation: Telethon Institute for Child Health Research under contract with the Department of Health, Western Australia, 2012.
43. Young J, Watson K, Ellis L, Raven L. Responding to the evidence: Breastfeed baby if you can – the sixth public health recommendation to reduce the risk of sudden unexpected death in infancy. *Breastfeeding Review* 2012; 20(1): 7-15.
44. Fetherston CM, Leach JL. Analysis of the ethical issues in the breastfeeding and bedsharing debate. *Breastfeeding Review* 2012; 20(3): 7-17.
45. Schluter PJ, Young J. Reducing the risk of Sudden Infant Death Syndrome: what infant care practices are being used by primary care-givers in Queensland? *Neonatal Paediatr Child Health Nurs* 2002; 5(2): 27-35.
46. Drever-Smith C, Bogossian F, New K. Co-sleeping and Bed sharing in Postnatal Maternity Units: A review of the literature and critique of clinical practice guidelines. *Int J Childbirth* 2013; 3(1): 13-27. <http://dx.doi.org/10.1891/2156-5287.3.1.13>

47. Royal College of Midwives. Bed-sharing and co-sleeping: Position Statement No 8. [Updated May 2004].
48. Australian Breastfeeding Association. Breastfeeding, Co-sleeping and Sudden Unexpected Deaths in Infancy. Authors: Kam R, Young J, 2013. Available from: [www.breastfeeding.asn.au/bfinfo/breastfeeding-co-sleeping-and-sudden-unexpected-deaths-infancy](http://www.breastfeeding.asn.au/bfinfo/breastfeeding-co-sleeping-and-sudden-unexpected-deaths-infancy).